

# MODELLING OF CIRCUMSTELLAR DUST AROUND RR TEL IN INFRARED

Tomislav Jurkić, Dubravka Kotnik-Karuza

Department of Physics, University of Rijeka, Croatia

## Observational data and methods of analysis

RR Tel JHKL magnitudes observed in the period 1975 - 2004 at SAAO have been analyzed. In order to show only the long-term variations, the **mira pulsation period** was determined and the light curves were **corrected for interstellar reddening and mira pulsations**. RR Tel gives evidence of **obscuration events** indicated in Fig. 1 (I, II and III).

The pulsation period was determined by fitting the JHKL light curves on a **periodic function** and in particular on a **sinusoid**, and by finally obtaining the residual light curve. The periodic function was fitted by dividing the light curve in bins for a chosen test period. By varying the number of bins and values of periods, the best fit was found using the least scattering method. Fitting on a sinusoid, the epochs of minima and maxima on the light curves were found by a fit on 3rd order Fourier polynomials, followed by determination of the initial pulsation epoch, pulsation period and amplitude.

By fitting JHKL magnitudes on 3rd order Fourier polynomials, we have **reconstructed the spectral energy distribution** (SED) of RR Tel at different time intervals when both JHKL magnitudes and ISO short wavelength spectra were available. The reconstructed SEDs cover the near- and mid-infrared spectral region in the periods with and without obscuration.

**Modeling** of circumstellar properties of the dust shell around the mira component was carried out by use of the numerical code **DUSTY**, assuming **spherical geometry**, with mira in the centre of the spherical dust shell. We used black body input radiation from the mira at a temperature between 2200 K and 2800 K. As the mira component has strong stellar winds, we assumed that envelope expansion is driven by radiation pressure on the dust grains. In the analytical approximation for **radiatively driven winds** the number density  $\eta$  is a function of the scaled radius  $y = r/r_{in}$ , of the initial  $v_i$  and final wind  $v_e$  velocity, while  $r_{in}$  is the inner dust shell radius (sublimation radius):

$$\eta \propto \frac{1}{y^2} \sqrt{\frac{y}{y-1 + (v_i/v_e)^2}}$$

For other forms of density distribution, the power law  $\eta \propto y^{-p}$  was used and power index  $p$  determined.

In the applied MRN dust grain size distribution  $n(a) \propto a^{-q}$ , ( $a_{min} < a < a_{max}$ ), the minimum grain size of  $a = 0.005 \mu m$  and  $q = 3.5$  are taken as fixed input parameters, while the maximum grain size  $a_{max}$  is a free parameter determined by modeling. The dust composition typical for mira stars containing **100% warm silicates** has been assumed. The outer dust shell radius is fixed to  $20 r_{in}$ , contrary to the inner dust shell radius  $r_{in}$  which is obtained by fitting, together with the dust sublimation temperature  $T_{dust}$

Symbiotic Mira	Other name	Spectral class		$A_v$	$d$ (kpc)	$P$ (days)	
RR Tel	Hen 3-1811	Mira M6/M5	Allen 1980, MNRAS, 192, 521 Muerseet & Schmid 1999, A&AS, 137, 473	0.3	2.5	387	Feast et al. 1983b, MNRAS, 202, 951 Penston et al. 1983, MNRAS, 202, 833 Kotnik-Karuza et al. 2006, A&A, 452, 503

## Results

### Pulsation period

	PERIODIC FUNCTION	SINUSOID	
	Period (days)	Period (days)	JD <sub>0</sub>
NO OBSCURATION	385.1 ± 4.8	384.3 ± 5.2	2 442 339 ± 7
OBSCURATION	386.4 ± 1.1	385.8 ± 2.3	2 442 344 ± 12
ALL INTERVALS	387.1 ± 0.8	386.8 ± 1.2	2 442 328 ± 11
Gromadzki et al, 2009		385 ± 4	2 442 303

The pulsation periods in epochs with and without obscuration have been determined using the two described methods and no statistically significant differences were found. This result suggests that there is no change in internal stellar properties which might cause obscuration influencing the dust properties. Periods obtained by two methods are almost undistinguishable, suggesting that the mira light curve variation is mainly sinusoidal. The fitted values agree very well with recent results from other authors, but are obtained with lower uncertainty.

### Conclusion

Single-shell dust model around mira with realistic grain sizes, chemical composition and radiatively driven winds can successfully explain near- and mid-infrared spectra of RR Tel symbiotic binary during obscuration events. Such events can be explained by dust condensed at distances beyond the inner dust shell radius and originating from the mira component. Modeling of circumstellar dust during obscuration events gives evidence of an increase in optical depth as well as in grain size from  $1.8 \mu m$  to  $4.0 \mu m$  as a consequence of possible grain growth, related to the unobscured epochs.

The SED modeling in the epoch without obscuration represents quite a challenge as both almost unattenuated stellar black body and strong dust emission are present, leading to a model of thick dust region of particular non-spherical configuration.

## References

Angeloni R. et al. 2010, MNRAS, 402, 2075  
Gromadzki M. et al. 2009, A&A, 59, 169  
Ivezic Z., Elitzur M. 1997, MNRAS, 287, 799  
Ivezic Z. et al. 1999, User Manual for DUSTY, University of Kentucky Internal Report, [www.pa.uky.edu/~moshe/dusty](http://www.pa.uky.edu/~moshe/dusty)  
Jurkić T., Kotnik-Karuza D. 2007, Balt.Astr., 16, 76  
Kotnik-Karuza D. et al. 2006, A&A, 452, 503  
Mathis J.S., Rumpel W., Nordsieck K.H. 1977, ApJ, 217, 425  
Mikolajewska J. 1999, in Optical and Infrared Spectroscopy of Circumstellar Matter, eds. E. W. Guenther, B. Stecklum, S. Klose, ASP Conf. Ser. 188, 291  
Ossenkopf V., Henning, Th., Mathis J.S. 1992, A&A, 261, 567

## Abstract

We present a circumstellar dust model around symbiotic mira RR Tel obtained by modelling the near-infrared JHKL magnitudes and ISO spectra. In order to follow the evolution of infrared colours in time, the published JHKL magnitudes were corrected by removing the mira pulsations. The RR Tel light curves show three obscuration events in the near-IR. Using the simultaneously available JHKL magnitudes and ISO spectra in three different epochs, we obtained SEDs in the near- and the mid-IR spectral region ( $1-20 \mu m$ ) in periods with and without obscuration. The pulsation periods determined from the JHKL photometry inside and outside obscuration intervals show no significant difference. The DUSTY numerical code was used to solve the radiative transfer and to determine the circumstellar dust properties of the inner dust regions around the mira, assuming spherical dust temperature distribution in its close neighbourhood. Dust temperature, grain size, density distribution, mass loss, terminal wind velocity and optical depth during intervals with and without obscuration have been obtained. Both JHKL and ISO observations during the obscuration period can be reproduced with the spherical dust envelope, while ISO spectra outside the obscuration show different behaviour. Possible configuration of the dust envelope has been discussed. Dynamical behaviour of the circumstellar dust was obtained by modelling the JHKL magnitudes observed during the span of more than 30 years.

## Results




Fig.1. JHKL lightcurves of RR Tel corrected for mira pulsations

## Dust properties




Fig.2. Two-colour diagram of RR Tel with theoretical models

RR Tel shows three distinct obscuration events. The single spherically-symmetric shell model with realistic physical dust properties can reproduce very well both long-term JHKL observations and ISO infrared spectra during these periods with obscuration.

According to our results, obscuration events can be explained by a change in dust optical depth, accompanied by an increase in dust grain size. Higher dust optical depth means larger amount of dust present around mira and hence, a higher mass loss.

An increase in dust grain size during obscuration can be caused by grain growth which agrees with an increase in mass loss as well as in optical depth.




Fig.3. ISO SWS spectra + SAAO JHKL during maximum obscuration (red arrow in Fig.1) (phase difference between epochs of two SEDs: 0.49 = 188 days)

Circumstellar dust modeling of the near-infrared JHKL magnitudes in epochs without obscuration results in a very thin dust shell and in low optical depth. The SED in the epoch without obscuration clearly shows almost unattenuated mira black body and strong dust emission. Such behaviour cannot be explained by a single dust shell, except in case of some other unknown mechanism or dust configuration which lead to a decrease of flux at wavelengths of about  $3 \mu m$ . In order to explain such behaviour, we have adopted a model in which mira is embedded in a thin dust shell and surrounded with thick probably toroidally configured emitting dust located outside our line of sight. Our results show that the dust region does not have a disk geometry, but is rather radially distributed by a steady-state wind, though radiatively driven winds cannot be discarded.




Fig.4. ISO SWS spectra + SAAO JHKL during minimum obscuration (blue arrow in Fig.1)

	OBSCURATION	NO OBSCURATION	
$T_{mira}$ (K)	2500 ± 300	2200	
		thin shell	'thick dust'
$T_{dust}$ (K)	1200 ± 100	1000 – 1200	1200
$a_{max}$ (μm)	4.0 ± 1.0	0.5 – 1.0	1.5 ± 0.5
$\tau_v$	5.2 – 5.8	0.4	12.5 ± 2.5
$A_K$	0.58 – 0.64	0.04	1.38 ± 0.28
$p$	rdw	rdw	2.0 or rdw
$\dot{M}$ ( $10^{-6} M_{\odot}/yr$ )	11 ± 1	0.4	
$v_e$ (km/s)	26 ± 2	28	

$T_{mira}$  mira temperature  
 $T_{dust}$  dust sublimation temperature  
 $a_{max}$  maximum grain size  
 $\tau_v$  visual optical depth  
 $A_K$  extinction at K  
 $p$  power index in density distribution  
 $\dot{M}$  mass-loss rate  
 $v_e$  terminal wind velocity  
rdw radiatively driven winds